

## ORIGINAL ARTICLE

# Dexmedetomidine or Propofol for Sedation in Mechanically Ventilated Adults with Sepsis

C.G. Hughes, P.T. Mailloux, J.W. Devlin, J.T. Swan, R.D. Sanders, A. Anzueto, J.C. Jackson, A.S. Hoskins, B.T. Pun, O.M. Orun, R. Raman, J.L. Stollings, A.L. Kiehl, M.S. Duprey, L.N. Bui, H.R. O Neal, Jr., A. Snyder, M.A. Gropper, K.K. Guntupalli, G.J. Stashenko, M.B. Patel, N.E. Brummel, T.D. Girard, R.S. Dittus, G.R. Bernard, E.W. Ely, and P.P. Pandharipande, for the MENDS2 Study Investigators\*

## ABSTRACT

**BACKGROUND**

Guidelines currently recommend targeting light sedation with dexmedetomidine or propofol for adults receiving mechanical ventilation. Differences exist between these sedatives in arousability, immunity, and inflammation. Whether they affect outcomes differentially in mechanically ventilated adults with sepsis undergoing light sedation is unknown.

**METHODS**

In a multicenter, double-blind trial, we randomly assigned mechanically ventilated adults with sepsis to receive dexmedetomidine (0.2 to 1.5  $\mu\text{g}$  per kilogram of body weight per hour) or propofol (5 to 50  $\mu\text{g}$  per kilogram per minute), with doses adjusted by bedside nurses to achieve target sedation goals set by clinicians according to the Richmond Agitation Sedation Scale (RASS, on which scores range from  $-5$  [unresponsive] to  $+4$  [combative]). The primary end point was days alive without delirium or coma during the 14-day intervention period. Secondary end points were ventilator-free days at 28 days, death at 90 days, and age-adjusted total score on the Telephone Interview for Cognitive Status questionnaire (TICS-T; scores range from 0 to 100, with a mean of  $50 \pm 10$  and lower scores indicating worse cognition) at 6 months.

**RESULTS**

Of 432 patients who underwent randomization, 422 were assigned to receive a trial drug and were included in the analyses. 214 patients received dexmedetomidine at a median dose of 0.27  $\mu\text{g}$  per kilogram per hour, and 208 received propofol at a median dose of 10.21  $\mu\text{g}$  per kilogram per minute. The median duration of receipt of the trial drugs was 3.0 days (interquartile range, 2.0 to 6.0), and the median RASS score was  $-2.0$  (interquartile range,  $-3.0$  to  $-1.0$ ). We found no difference between dexmedetomidine and propofol in the number of days alive without delirium or coma (adjusted median, 10.7 vs. 10.8 days; odds ratio, 0.96; 95% confidence interval [CI], 0.74 to 1.26), ventilator-free days (adjusted median, 23.7 vs. 24.0 days; odds ratio, 0.98; 95% CI, 0.63 to 1.51), death at 90 days (38% vs. 39%; hazard ratio, 1.06; 95% CI, 0.74 to 1.52), or TICS-T score at 6 months (adjusted median score, 40.9 vs. 41.4; odds ratio, 0.94; 95% CI, 0.66 to 1.33). Safety end points were similar in the two groups.

**CONCLUSIONS**

Among mechanically ventilated adults with sepsis who were being treated with recommended light-sedation approaches, outcomes in patients who received dexmedetomidine did not differ from outcomes in those who received propofol. (Funded by the National Institutes of Health; ClinicalTrials.gov number, NCT01739933.)

The authors' full names, academic degrees, and affiliations are listed in the Appendix. Address reprint requests to Dr. Hughes at 1211 21st Ave. S., 422 MAB, Nashville, TN 37212, or at christopher.hughes@vumc.org.

\*The MENDS2 Study Investigators are listed in the Supplementary Appendix, available at NEJM.org.

This article was published on February 2, 2021, at NEJM.org.

N Engl J Med 2021;384:1424-36.

DOI: 10.1056/NEJMoa2024922

Copyright © 2021 Massachusetts Medical Society.

WORLDWIDE, AT LEAST 20 MILLION patients each year have sepsis with severe organ dysfunction,<sup>1</sup> with over 20% receiving mechanical ventilation.<sup>2,3</sup> Sedative medications are frequently used for patient comfort and safety but may potentiate acute brain dysfunction (e.g., delirium or coma) and long-term cognitive impairment.<sup>4-10</sup> Basic and translational studies show that among the recommended sedatives, dexmedetomidine (an  $\alpha_2$  receptor agonist) has antiinflammatory and bacterial clearance properties that are superior to those of gamma-aminobutyric acid (GABA) agonists, such as benzodiazepines and propofol, and also reduces neuronal apoptosis and promotes biomimetic sleep—all of which could improve clinical outcomes.<sup>11-17</sup> Trials comparing dexmedetomidine with benzodiazepines in adults have shown that the use of dexmedetomidine results in improvement in outcomes such as delirium, coma, and time receiving mechanical ventilation.<sup>18,19</sup> Patients treated with dexmedetomidine had a lower incidence of subsequent infection,<sup>19</sup> and the beneficial effects of dexmedetomidine, including lower 28-day mortality, were more pronounced in patients with sepsis.<sup>18,20</sup>

A noninferiority trial comparing dexmedetomidine with propofol in critically ill patients, about half of whom had sepsis, showed that patients who received dexmedetomidine were more interactive, but the choice of sedation did not affect the duration of mechanical ventilation, the length of stay in the intensive care unit (ICU) or hospital, or short-term mortality.<sup>21</sup> The differences between the sedatives with respect to the risk of acute brain dysfunction or cognitive impairment and mortality months after critical illness were unclear. Subsequent open-label trials with dexmedetomidine as the primary sedative did not show a reduction in acute brain dysfunction, a greater number of ventilator-free days, or lower mortality at 180 days than was shown with control sedative regimens (primarily propofol), although concomitant nontrial sedatives were frequently used and few patients were maintained at light sedation.<sup>22,23</sup>

The Society of Critical Care Medicine<sup>24</sup> recommends sedation with either dexmedetomidine or propofol targeted to light levels of sedation for adults receiving mechanical ventilation and continuous sedation. Given the superior immunomodulatory effects of dexmedetomidine and its benefit in patients with sepsis as compared with

benzodiazepines, we designed the MENDS2 trial (Maximizing the Efficacy of Sedation and Reducing Neurological Dysfunction and Mortality in Septic Patients with Acute Respiratory Failure) to test whether dexmedetomidine leads to better short-term and long-term outcomes than propofol in mechanically ventilated adults with sepsis.

## METHODS

### TRIAL DESIGN AND OVERSIGHT

We conducted a double-blind, randomized, controlled trial at 13 medical centers in the United States. The institutional review board at each center approved the protocol (available with the full text of this article at NEJM.org). Patients or their surrogates provided written informed consent before enrollment. The trial was designed by the authors, who gathered and analyzed the data, attest to the accuracy and completeness of the data, vouch for the fidelity of the trial to the protocol, and wrote and agreed to submit the manuscript for publication. An independent data and safety monitoring board provided oversight of the trial. Pfizer supplied the dexmedetomidine trial drug but had no role in the design or conduct of the trial, analysis of the data, or writing of the manuscript. The Food and Drug Administration approved an Investigational New Drug application for dexmedetomidine administered for more than 24 hours and for doses up to 1.5  $\mu\text{g}$  per kilogram per hour (see the Supplementary Appendix, available at NEJM.org). We registered the trial at ClinicalTrials.gov before enrollment began. Before group assignments were unmasked, we registered the statistical analysis plan at Open Science Framework (<https://osf.io/dfyxh/>) in January 2019 (with publication in March 2020).<sup>25</sup>

### PATIENT SELECTION AND RANDOMIZATION

We included adults who were sequentially admitted to a medical or surgical ICU, had suspected or known infection, and were treated with continuous sedation for invasive mechanical ventilation. Patients were excluded if they had baseline severe cognitive impairment; were pregnant or breast-feeding; were blind, deaf, or unable to understand approved languages; had second-degree or third-degree heart block or persistent bradycardia requiring intervention; had an allergy to dexmedetomidine or propofol; had an indication for benzodiazepines; were anticipated to have immediate discontinuation of mechanical ventila-

tion; were expected to have neuromuscular blockade for more than 48 hours; were in a moribund state; or had received mechanical ventilation for more than 96 hours before meeting all inclusion criteria. Additional details on exclusion and inclusion criteria are provided in Section S1 in the Supplementary Appendix. We randomly assigned patients to receive dexmedetomidine or propofol in a 1:1 ratio using computer-generated permuted blocks stratified by enrollment site and age (<65 years vs. ≥65 years). Researchers, clinicians (except bedside nurses), patients, and families were unaware of the group assignments.

#### TRIAL INTERVENTIONS AND MEASUREMENTS

Investigational pharmacists prepared dexmedetomidine (5 µg per milliliter) and propofol (10 mg per milliliter) in identical intravenous fluid bags covered with opaque plastic bags to be administered in units of milliliters per hour to maintain study masking (Sections S2 and S3). Bedside nurses covered intravenous tubing with opaque coverings and verified that covers were in place before study personnel or clinicians entered patients' rooms. The trial drug was initially infused at a dose corresponding to the same sedative dosing that the patient was receiving immediately before randomization. Bedside nurses used a weight-based dosing guideline (0.15 to 1.5 µg per kilogram of actual body weight per hour for dexmedetomidine and 5 to 50 µg per kilogram of actual body weight per minute for propofol) to adjust the trial drug every 10 minutes to target sedation goals set by the clinical team and documented each adjustment and the rationale for it. The clinical team used the Richmond Agitation Sedation Scale (RASS, on which scores range from −5 [unresponsive] to +4 [combative]),<sup>26</sup> to set the sedation goal, which was primarily light sedation (RASS score 0 to −2).

Administration of the trial drug was temporarily held in the event of hypotension, bradycardia, sedation deeper than the target level, spontaneous awakening trials, or surgery. The trial drug was permanently discontinued if the patient had persistent symptomatic bradycardia, new onset second- or third-degree heart block, serious allergic reactions, suspected propofol-related infusion syndrome (refractory shock, rhabdomyolysis, acidosis, and kidney failure related to high propofol exposure), or any serious adverse event related to the intervention. The trial drug was discontinued

after the 14-day intervention period, extubation, or discharge from the ICU, whichever came first. Patients whose trachea was extubated and reintubated within the 14-day intervention period resumed the trial drug if sedation was indicated.

We treated pain with intermittent opioid boluses or fentanyl infusion (see Section S4 for details regarding rescue sedation, neuromuscular blockade, and treatment of agitated delirium). Additional patient care practices (e.g., administration of fluids, vasopressors, or antibiotics and extubation criteria) were based on international guideline recommendations.<sup>24,27</sup>

All centers performed, and investigators reinforced, the ABCDE (awakening and breathing coordination, choice of sedation, delirium monitoring and management, and early mobility) bundle,<sup>28,29</sup> with daily adherence recorded. In addition to care assessments made by nurses, trained research personnel assessed patients with the use of RASS for level of arousal,<sup>26</sup> Confusion Assessment Method for the ICU (CAM-ICU)<sup>30</sup> for delirium, and the Critical Care Pain Observation Tool<sup>31</sup> for pain; assessments were made twice daily in the ICU and then once daily after transfer from the ICU for up to 14 days or until discharge from the hospital or death. We strived to conduct delirium assessments when the patient was maximally awake. A RASS score of −4 or −5 indicated coma, and a positive CAM-ICU score indicated delirium.

Six months after randomization, research personnel assessed patients' cognition with the Telephone Interview for Cognitive Status (TICS) questionnaire<sup>32</sup> and a validated telephone cognitive battery,<sup>33</sup> functional status with the Katz Activities of Daily Living (ADL) scale<sup>34</sup> and the Functional Activities Questionnaire (FAQ),<sup>35</sup> and quality of life with the European Quality of Life 5 Dimensions (EQ-5D) survey (Section S5).<sup>36</sup>

#### TRIAL END POINTS

The primary efficacy end point was the number of calendar days alive without delirium or coma during the 14-day intervention period. Secondary efficacy end points included ventilator-free days at 28 days, death at 90 days, and global cognition at 6 months using the age-adjusted TICS total score (TICS-T score). Additional details regarding efficacy, adherence, and safety end points are provided in Section S5, along with a list of additional end points not reported here.

**STATISTICAL ANALYSIS**

Owing to enrollment that was slower than anticipated, the data and safety monitoring board and the National Institutes of Health approved a protocol amendment in March 2017 to lower the enrollment target from 530 patients to 420 patients receiving the trial drug to provide 85% power to detect a 1.5-day difference in days alive without delirium or coma between groups and 80% power to detect a 12 percentage-point absolute difference in mortality at 90 days, assuming an expected mortality of 30% in the propofol group. We had at least 80% power to detect a 3.9-point difference in age-adjusted TICS-T scores between groups, with a 5-point difference considered to be clinically important.

We analyzed data in the modified intention-to-treat population, which was prespecified as all patients who underwent randomization and received a trial drug. We analyzed primary and secondary end points using both univariate methods and multivariable regression models and considered adjusted analyses to be the primary analyses. We analyzed days alive without delirium and coma, ventilator-free days, and age-adjusted TICS-T scores at 6 months using proportional-odds logistic regression and analyzed death at 90 days using Cox proportional-hazards regression (adjusted for covariates listed in Section S5).

We adjusted the level of statistical significance for the primary end point analysis to  $P < 0.044$  to account for one prespecified planned interim analysis. The level of statistical significance for all other end points was  $P < 0.05$ . Simple imputation was used for missing in-hospital variables and multiple imputation for partially available long-term end points to avoid bias owing to missing variables. We did not adjust for multiple comparisons in the analysis of secondary end points. We used Research Electronic Data Capture software (REDCap, Vanderbilt University) for data management and R, version 3.6.2 (R Foundation for Statistical Computing), for statistical analyses.


**RESULTS****PATIENTS**

From May 2013 through December 2018, we screened 4840 patients, 4402 (91%) of whom met at least one exclusion criterion (Fig. 1). Of 438 patients enrolled, 6 were subsequently found to

be ineligible, 432 patients underwent randomization, and 422 began receiving dexmedetomidine (214 patients) or propofol (208 patients). The demographic and in-hospital characteristics of the patients are shown in Table 1 and Table S1.

**TRIAL INTERVENTIONS**

Details of trial drug dosing, dose adjustment, and sedation regimen are shown in Table 2 and Table S2. The median RASS score as assessed by the research team was  $-2$  (interquartile range,  $-3.00$  to  $-1.00$ ) while patients were receiving a trial drug (median days of administration, 3.0 [interquartile range, 2.0 to 6.0]), indicating light sedation and ability of patients to make eye contact with only verbal stimulation (Fig. S1). The overall time spent at the target sedation was close to 60% in both groups (Fig. S2). Bedside nurses adjusted the trial drug infusion a median of 10 times (interquartile range, 5 to 21) over the duration of administration. Common reasons for adjustment of the infusion included undersedation, oversedation, and hypotension. The trial drug was temporarily held in approximately one quarter of all patients. Rescue midazolam was used in about half the patients, most often for procedural sedation or during neuromuscular blockade, and the median daily exposure on days it was administered was 4 mg (interquartile range, 2 to 11). The use of open-label propofol (received by 13% in the dexmedetomidine group and 8% in the propofol group) and dexmedetomidine (4% in the dexmedetomidine group and 3% in the propofol group) was infrequent and doses were low, indicating high adherence to the protocol. Overall, 42% of the patients received an antipsychotic medication. Soft wrist restraints during receipt of mechanical ventilation were the standard of care in our trial ICUs; thus, 96% of patients had restraints in place for a median of 3 days (interquartile range, 2 to 5). Neuromuscular blockade infusion was used in 17% of patients for a median of 1 day (interquartile range, 1 to 2) at some point while they were receiving the trial drug. Pain was well controlled in both groups according to Critical Care Pain Observation Tool scoring, and we noted high adherence to all components of the ABCDE bundle. We proactively assessed for unblinding among clinicians and research staff and found an episode of unblinding in 58 patients (14%), with a similar frequency in the two groups.





**Figure 1 (facing page). Screening, Randomization, Follow-up, and Analysis.**

The number of patients excluded for each criterion total more than the total number of patients excluded because some patients met more than one exclusion criterion.

**EFFICACY END POINTS**

The adjusted number of days alive without delirium or coma over the 14-day intervention period was not significantly different between the dexmedetomidine group (adjusted median, 10.7 days; 95% confidence interval [CI], 8.5 to 12.5) and the propofol group (adjusted median, 10.8 days; 95% CI, 8.7 to 12.6) (odds ratio, 0.96; 95% CI, 0.74 to 1.26;  $P=0.79$ ). Similarly, we found no significant differences between the dexmedetomidine and propofol groups in the number of ventilator-free days at 28 days (adjusted median, 23.7 vs. 24.0 days; odds ratio, 0.98; 95% CI, 0.63 to 1.51) or in death at 90 days (81 patients [38%] vs. 82 patients [39%]; hazard ratio, 1.06; 95% CI, 0.74 to 1.52). Results of primary and secondary efficacy end point analyses are shown in Table 3, Figure 2, and Figure S3.

We assessed more than 90% of eligible patients at 6 months after randomization (Fig. 1). Approximately 25% in each group had age-adjusted TICS-T scores that were 2 standard deviations below population norms (i.e., a score of  $\leq 30$ , at a level consistent with impairment), which suggests clinically important cognitive dysfunction 6 months after critical illness. We observed no significant differences between the dexmedetomidine and propofol groups in age-adjusted TICS-T scores at 6 months (adjusted median score, 40.9 vs. 41.4; odds ratio, 0.94; 95% CI, 0.66 to 1.33). There were no clinically meaningful differences between groups in median cognitive, functional, and quality-of-life assessment scores at 6 months (Table S4).

Results of sensitivity analyses that included the 10 patients who underwent randomization but never received a trial drug (Table S5) were qualitatively similar to the results of the main analyses. Results of differential effects of the study treatment on end points according to age at enrollment, baseline cognition, and medical as compared with surgical hospitalization show that the clinical importance of these interactions appeared to be minimal (Figs. S4 through S9); however,

the trial may not have been adequately powered to draw conclusions about these or other subgroups.

**SAFETY END POINTS**

Data on organ dysfunction and safety end points by group are shown in Tables S6 and S7, respectively. The proportions of patients who had organ dysfunction, hypotension, or severe lactic acidosis after randomization were similar in the two groups. Symptomatic bradycardia requiring discontinuation of the trial drug was similar in the two groups (Table S2). Fewer patients in the dexmedetomidine group had acute respiratory distress syndrome (ARDS) or signs of trial drug withdrawal, and fewer patients in the propofol group extubated themselves. Median plasma triglyceride levels and the proportion of patients with severely elevated levels of triglyceride ( $>500$  mg per deciliter) were quantitatively higher in the propofol group than in the dexmedetomidine group on days 7 and 14, although these differences are unlikely to be clinically relevant. Similarly, median plasma cortisol levels at day 14 were slightly lower in the dexmedetomidine group than in the propofol group, including a higher proportion of patients with low cortisol ( $<20$   $\mu\text{g}$  per deciliter). Clinicians had access to these results without indication of group assignment and discontinued the trial drug in eight patients owing to hypertriglyceridemia. One patient had suspected propofol-related infusion syndrome (later disproved) and had the propofol discontinued.

**DISCUSSION**

In this multicenter, double-blind, randomized, controlled trial involving mechanically ventilated adults with sepsis who were being treated with recommended light-sedation approaches, we did not find evidence that sedation with dexmedetomidine led to more days alive without acute brain dysfunction than propofol. Furthermore, we found no difference in ventilator-free days at 28 days, death at 90 days, or global cognition (as assessed with the use of age-adjusted TICS-T scores) at 6 months between the dexmedetomidine and propofol groups. Safety end points were also similar in the two groups.

Although recent data suggest that many criti-

**Table 1. Baseline Characteristics of the Patients.\***

Characteristic	Dexmedetomidine (N=214)	Propofol (N=208)
Median age (IQR) yr	59 (48 68)	60 (50 68)
Female sex no. (%)	93 (43)	88 (42)
Median body-mass index (IQR)	30 (25 38)	29 (25 37)
Race or ethnic group no. (%)		
White	188 (88)	177 (85)
Black	15 (7)	23 (11)
Latinx	12 (6)	18 (9)
Multiple or other	11 (5)	8 (4)
Median IQCODE-SF score (IQR)	3.06 (3.00 3.23)	3.00 (3.00 3.25)
Median Charlson Comorbidities Index score (IQR)	2 (1 4)	2 (1 4)
Admitted to surgical ICU no. (%)	76 (36)	72 (35)
Median APACHE II score at ICU admission (IQR)	27 (21 32)	27 (22 32)
Median days from ICU admission to trial enrollment (IQR)	1.21 (0.67 1.95)	1.17 (0.68 1.94)
Median days of mechanical ventilation before trial enrollment (IQR)	0.98 (0.58 1.36)	0.97 (0.61 1.54)
Median total SOFA score at trial enrollment (IQR)**	10 (8 13)	10 (8 12)
Shock, receiving vasopressor, at enrollment no. (%)	119 (56)	102 (49)
Known or suspected source of infection no. (%)		
Blood	92 (43)	79 (38)
Lung	116 (54)	133 (64)
Abdomen	19 (9)	20 (10)
Urinary tract	46 (21)	55 (26)
Skin or wound	23 (11)	26 (12)
Stool	12 (6)	12 (6)
Other	24 (11)	21 (10)
Infection status no. (%)		
Infection confirmed by culture	146 (68)	132 (63)
Infection suspected but not confirmed by culture	58 (27)	68 (33)
Infection ruled out	10 (5)	8 (4)
Dexmedetomidine before enrollment no. (%)	35 (16)	25 (12)
Propofol before enrollment no. (%)	131 (61)	129 (62)
Benzodiazepine before enrollment no. (%)	62 (29)	73 (35)
Opioid before enrollment no. (%)	144 (67)	147 (71)
Antipsychotic agent before enrollment no. (%)	24 (11)	27 (13)
Delirium at enrollment no. (%)	75 (35)	91 (44)
Level of arousal closest to the time of randomization no. (%)		
Coma: RASS -5 or -4	81 (38)	74 (36)
Deep sedation: RASS -3	29 (14)	38 (18)
Light sedation: RASS -2 or -1	85 (40)	75 (36)
Awake and calm: RASS 0	13 (6)	14 (7)
Agitated: RASS +1 to +4	6 (3)	7 (3)

\* Percentages may not total 100 because of rounding. Summary statistics are reported for nonmissing values. ICU denotes intensive care unit, and IQR interquartile range.

The body-mass index is the weight in kilograms divided by the square of the height in meters.

Race or ethnic group was reported by the patient or determined by the treating physicians.

The Informant Questionnaire on Cognitive Decline in the Elderly short form (IQCODE-SF)<sup>37</sup> was used to determine preexisting dementia; scores range from 1.0 to 5.0, with higher scores indicating more severe cognitive impairment. Scores on the Charlson Comorbidity Index range from 0 to 33, with higher scores indicating a higher risk of death from a coexisting illness.

|| The Acute Physiology and Chronic Health Evaluation (APACHE II) assesses the risk of death on a scale from 0 to 71, with higher scores indicating a higher risk of death.

\*\* The Sequential Organ Failure Assessment (SOFA) is used to track organ failure in the ICU; scores range from 0 to 24, with higher scores indicating greater severity of illness.

Delirium was deemed to be present when the Confusion Assessment Method for the Intensive Care Unit (CAM-ICU, which scores delirium as either present [positive] or not present [negative]), was positive.

The Richmond Agitation Sedation Scale (RASS) measures levels of consciousness on a scale from -5 (unresponsive) to +4 (combative).

**Table 2. Adherence and Sedation Regimen.**

Outcome	Dexmedetomidine N=214	Propofol N=208
Median hours from meeting inclusion criteria to drug initiation (IQR)	22.4 (13.4 31.3)	22.1 (12.8 33.7)
Median hours from randomization to drug initiation (IQR)	1.3 (0.9 2.2)	1.3 (0.8 2.1)
Trial drug administration		
Median days of receipt of drug (IQR)	3.0 (2.0 5.0)	4.0 (2.0 6.0)
Median days from first meeting trial criteria to initiation of drug (IQR)	1.00 (0.00 1.00)	1.00 (0.00 1.00)
Median daily volume on days administered (IQR) ml	119 (46 243)	131 (67 229)
Median daily dose on days administered (IQR)	0.27 $\mu$ g/kg/hr (0.11 0.61)	10.2 $\mu$ g/kg/min (5.5 18.4)
Median total no. of drug adjustments per patient (IQR)	9 (5 15.8)	11.5 (5.8 25)
Drug temporarily held no. (%) <sup>*</sup>	60 (28)	57 (27)
Median no. of times drug temporarily held per patient (IQR)	1 (1 1)	1 (1 2)
Drug permanently discontinued no. (%)	25 (12)	23 (11)
Trial or clinical team aware of the drug used no. (%)	27 (13)	31 (15)
Withdrawal from trial during hospitalization no. (%)	10 (5)	9 (4)
Median RASS score while receiving drug (IQR)	-2.00 (-3.00 to -1.00)	-1.95 (-3.03 to -0.98)
Percent time at target sedation level while receiving drug	57	60
Median CPOT score while receiving drug (IQR)	0.33 (0.00 0.83)	0.31 (0.00 0.87)
Percent of days with adherence to ABCDE bundle		
Spontaneous awakening trial	98	98
Spontaneous breathing trial	93	95
Coordination of awakening and breathing trials	86	84
Nondrug delirium interventions	99	99
Early mobilization	91	92
Median daily fentanyl dose on days administered (IQR) $\mu$ g/hr	68 (28 119)	56 (20 95)
Midazolam exposure		
Ever used no. (%)	114 (53)	90 (43)
Median days among users (IQR)	2.0 (1.0 4.0)	1.0 (1.0 2.0)
Median daily dose on days administered (IQR) mg per day	3.8 (2.0 10.9)	4.0 (2.0 10.8)
Antipsychotic exposure		
Ever used no. (%)	90 (42)	87 (42)
Median days among users (IQR)	5.0 (2.0 7.8)	4.0 (2.0 8.0)
Median daily dose on days administered (IQR) mg	2.2 (1.0 6.4)	3.6 (1.0 6.3)
Open-label propofol exposure		
Ever used no. (%)	27 (13)	16 (8)
Median days among users (IQR)	2.0 (1.0 3.0)	1.5 (1.0 2.0)
Median daily dose on days administered (IQR) $\mu$ g/kg/min	10.8 (4.9 17.4)	4.8 (3.4 6.6)
Open-label dexmedetomidine exposure		
Ever used no. (%)	9 (4)	6 (3)
Median days among users (IQR)	1.0 (1.0 2.0)	1.0 (1.0 3.2)
Median daily dose on days administered (IQR) $\mu$ g/kg/hr	0.24 (0.04 0.30)	0.26 (0.07 0.7)

\* The reasons for temporary holding of the drug included oversedation, hypotension, or bradycardia; spontaneous awakening trials or times during which patients were not being sedated, were not receiving mechanical ventilation, or were in the operating room are not included.

The Critical Care Pain Observation Tool (CPOT) is used to assess for pain by evaluating facial expression, body movement, muscle tension, and adherence to use of the ventilator if intubated or vocalization if extubated. Total scores range from 0 to 8, with scores higher than 2 indicating the presence of pain.

The ABCDE bundle includes evaluations for awakening and breathing coordination, choice of sedation, delirium monitoring and management, and early mobility.

Values shown are in intravenous haloperidol equivalents.



**Table 3. Primary and Secondary Efficacy End Points.\***

End Point	Dexmedetomidine (N=214)	Propofol (N=208)
<b>Primary end point</b>		
Days alive without delirium or coma at 14 days		
Unadjusted no. of days    median (IQR)	8.0 (1.0 12.8)	7.5 (1.8 11.2)
Adjusted no. of days    median (95% CI)	10.7 (8.5 12.5)	10.8 (8.7 12.6)
Adjusted odds ratio (95% CI)	0.96 (0.74 1.26)	Reference
<b>Secondary end points</b>		
Ventilator-free days at 28 days		
Unadjusted no. of days    median (IQR)	20.9 (0.0 26.1)	19.9 (4.2 24.9)
Adjusted no. of days    median (95% CI)	23.7 (20.5 25.4)	24.0 (20.9 25.4)
Adjusted odds ratio (95% CI)	0.98 (0.63 1.51)	Reference
Death at 90 days		
Unadjusted no. of patients (%)	81 (38)	82 (39)
Adjusted hazard ratio (95% CI)	1.06 (0.74 1.52)	Reference
TICS-T score at 6 mo		
Unadjusted score    median (IQR)	39 (28 48)	38 (30 46)
Adjusted score    median (95% CI)	40.9 (33.6 47.1)	41.4 (34.0 47.3)
Adjusted odds ratio (95% CI)	0.94 (0.66 1.33)	Reference


\* Variables in adjusted analyses, except for analysis of death at 90 days, included the following: age at trial enrollment; education; baseline cognitive function as determined according to the IQCODE-SF; preexisting coexisting conditions according to the Charlson Comorbidities Index; SOFA assessment on the day of enrollment (excluding central nervous system component); level of arousal at randomization according to the RASS score closest to the time of randomization; exposure to propofol, dexmedetomidine, benzodiazepines, opioids, and antipsychotics between the time of ICU admission and midnight before enrollment; medical (vs. surgical) patient; and infection type. Variables in adjusted analyses for death at 90 days included the following: age at trial enrollment, baseline cognitive function as determined according to the IQCODE-SF, preexisting coexisting conditions according to the Charlson Comorbidities Index, SOFA assessment on the day of enrollment (excluding central nervous system component), medical (vs. surgical) patient, and infection type.

Age-adjusted total scores on the Telephone Interview for Cognitive Status questionnaire (TICS-T) range from 0 to 100 with a mean of 50±10; lower scores indicate worse cognition, and a score of 35 or less indicates cognitive impairment.

cally ill adults receiving mechanical ventilation may not require sedative infusions,<sup>38,39</sup> our trial specifically enrolled adults with sepsis who had a high severity of illness, a greater risk for ARDS, and a higher requirement for continuous sedation. It was important to better characterize the effect of greater arousability, analgesic properties, and lack of respiratory depression observed with dexmedetomidine as compared with GABAergic sedatives in this population. Data indicate meaningful differences between dexmedetomidine and GABAergic sedatives with respect to innate immunity and risk of infection, including evidence that dexmedetomidine may offer superior anti-inflammatory effects.<sup>11-17</sup> Despite these theoretical benefits and studies supporting the use of

dexmedetomidine, the choice between dexmedetomidine and propofol alone does not appear to substantially affect patient outcomes in the complex milieu of critical illness with sepsis. Our findings, therefore, strongly reinforce current guidelines<sup>24</sup> that recommend the use of either dexmedetomidine or propofol for light sedation when continuous sedation is needed for adults with or without sepsis who require mechanical ventilation.

Our trial builds on other trials that have compared dexmedetomidine with propofol,<sup>21-23,40</sup> with important methodologic advances that include a higher degree of sedative trial drug blinding, a better separation between groups with regard to sedative exposure, and stricter adherence to light



sedation approaches, with high compliance with a standardized, multicomponent sedation management bundle (i.e., the ABCDE bundle)<sup>28,29</sup> that has been shown to reduce mortality and improve other important outcomes. One study by Kawazoe et al.<sup>22</sup> randomly assigned 201 patients with sepsis who required mechanical ventilation to open-label sedation with dexmedetomidine (up to 0.7  $\mu\text{g}$  per kilogram per hour) or sedation without dexmedetomidine (infusions of propofol or midazolam or both) for up to 7 days. On 1 or more study days, 29% of the dexmedetomidine group received propofol (nearly three times the crossover rate of our study) and up to 21% received midazolam. The authors found no significant difference in the number of days without delirium or coma, the number of ventilator-free days, or mortality at 28 days with dexmedetomidine use, although the trial was probably underpowered to measure a difference in mortality.

More recently, Shehabi et al.<sup>23</sup> performed a landmark open-label, randomized trial of dexmedetomidine (up to 1.5  $\mu\text{g}$  per kilogram per hour) as compared with usual care (infusions of propofol

or midazolam or both) for up to 28 days in more than 3900 patients with critical illness. The authors did not find a significant difference between the groups in the number of days without delirium or coma at 28 days, the number of ventilator-free days at 28 days, death at 90 days (including in subgroup analyses of 806 patients with suspected or confirmed sepsis), or death at 180 days. Most patients (86%) in the dexmedetomidine group received concomitant propofol for a median of 2.0 days, and 23% received midazolam for a median of 0.5 days; this lack of separation between groups limits the interpretation of the results. Despite an unmasking episode in 14% of patients and crossover in about 10% of patients in the present study, we believe that our methodologic rigor allows a more definitive conclusion that dexmedetomidine and propofol have similar efficacy with regard to acute brain dysfunction, mechanical ventilation requirement, and mortality when light sedation goals and the ABCDE bundle are used to care for critically ill mechanically ventilated adults with sepsis. Biologically, patients with sepsis should derive im-

portant benefits from dexmedetomidine because of its immunomodulatory and antiinflammatory properties; thus, it is highly unlikely that patients without sepsis would have outcomes with dexmedetomidine substantially different from those we report.

An expanding area of interest in the care of critically ill patients is the prevention of cognitive impairment, functional impairment, and decline in quality of life after hospital discharge. The study by Shehabi et al.<sup>23</sup> showed similar scores in cognition (as assessed with the Informant Questionnaire on Cognitive Decline in the Elderly [IQCODE]) and quality of life (as assessed with the EQ-5D) at 180 days in the dexmedetomidine and control groups. Using a more robust cognitive assessment battery, we found clinically important cognitive dysfunction in approximately 25% of patients after sepsis and critical illness even with light sedation approaches, and the use of dexmedetomidine as compared with propofol did not alter this finding. Considering our high follow-up rates and use of a robust assessment battery, it appears that sedation choice does not affect survivorship outcomes when currently recommended sedation approaches are used.

Our trial has a number of strengths but also some notable limitations. We made every effort to mask the delivery of propofol and dexmedetomidine considering their different physical properties. Although an episode of unmasking of the group assignment to a clinician or research team member occurred in 14% of patients, adherence to blinding in our trial was higher than that reported in similar clinical trials of propofol and dexmedetomidine. We allowed clinicians to set sedation targets, achieved good separation between groups regarding sedative exposure, and had robust follow-up. In general, patients had light levels of sedation with low doses of sedative medications and concomitant opioid analgesia.

This may reflect changing sedation strategies conforming to recommended practices or the need for lower sedative doses in patients with sepsis. We had some cross-contamination of sedative use, although substantially less than that in similar sedation studies, and had a rescue protocol that included the use of low-dose antipsychotic medications. The trial drug was started a median of 22 hours after the patient met all inclusion criteria, which may have limited our ability to affect outcomes. We had slower-than-anticipated enrollment, which required an adjustment of the sample size, yet had adequate power to study the questions of interest. Some exclusions were the result of clinicians not having equipoise regarding sedation for a given patient or were due to patients (or their surrogates) decision not to agree to enrollment in the trial, factors that may affect generalizability. Overall, we believe that we studied a representative population of patients with sepsis in centers across the United States and provide more definitive evidence regarding the choice of sedation in critically ill patients with sepsis who require mechanical ventilation.

Our trial showed that among critically ill adults with sepsis who were receiving mechanical ventilation and for whom recommended light-sedation approaches were used, dexmedetomidine did not lead to better outcomes than propofol with respect to days alive without acute brain dysfunction, ventilator-free days, death at 90 days, or cognition at 6 months.

Supported by a grant (HL111111) from the National Institutes of Health (NIH), a grant (TR000445) from the NIH for the use of REDCap, grants from the NIH to Dr. Duprey (1F31AG066460) and to Dr. Girard (HL135144), and a grant from the National Heart Lung and Blood Institute to Dr. Pun (HL14678-01). Dexmedetomidine was provided by Pfizer (previously by Hospira).

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

A data sharing statement provided by the authors is available with the full text of this article at NEJM.org.

#### APPENDIX

The authors' full names and academic degrees are as follows: Christopher G. Hughes, M.D., M.S.C.R., Patrick T. Mailloux, D.O., John W. Devlin, Pharm.D., Joshua T. Swan, Pharm.D., M.P.H., Robert D. Sanders, M.D., Antonio Anzueto, M.D., James C. Jackson, Psy.D., Aimee S. Hoskins, B.S.N., R.N., Brenda T. Pun, D.N.P., R.N., Onur M. Orun, M.S., Rameela Raman, Ph.D., Joanna L. Stollings, Pharm.D., Amy L. Kiehl, M.A., Matthew S. Duprey, Pharm.D., Ph.D., Lan N. Bui, Pharm.D., M.P.H., Hollis R. O'Neal, Jr., M.D., Allison Snyder, M.S.N., A.P.R.N., Michael A. Gropper, M.D., Ph.D., Kalpalatha K. Guntupalli, M.D., Gregg J. Stashenko, M.D., Mayur B. Patel, M.D., M.P.H., Nathan E. Brummel, M.D., M.S.C.I., Timothy D. Girard, M.D., M.S.C.I., Robert S. Dittus, M.D., M.P.H., Gordon R. Bernard, M.D., E. Wesley Ely, M.D., M.P.H., and Pratik P. Pandharipande, M.D., M.S.C.I.

The authors' affiliations are as follows: the Critical Illness, Brain Dysfunction, and Survivorship Center (C.G.H., J.C.J., A.S.H., B.T.P., O.M.O., R.R., J.L.S., A.L.K., M.B.P., N.E.B., T.D.G., R.S.D., G.R.B., E.W.E., P.P.P.), the Center for Health Services Research (C.G.H., J.C.J., R.R., M.B.P., T.D.G., R.S.D., E.W.E., P.P.P.), the Division of Anesthesiology Critical Care Medicine, Department of Anesthesiol-

ogy (C.G.H., P.P.P.), the Division of Allergy, Pulmonary, and Critical Care Medicine (J.C.J., B.T.P., G.R.B., E.W.E.), and the Division of General Internal Medicine and Public Health (R.S.D.), Department of Medicine, the Departments of Biostatistics (O.M.O., R.R.) and Pharmaceutical Services (J.L.S.), and Division of Trauma and Surgical Critical Care, Department of Surgery (M.B.P.), Vanderbilt University Medical Center, and the Anesthesia Service (C.G.H., P.P.P.), Research Service (J.C.J.), Surgical Service (M.B.P.), and Geriatric Research, Education and Clinical Center (R.S.D., E.W.E.), Department of Veterans Affairs Medical Center, Tennessee Valley Healthcare System both in Nashville; the Neuroscience Institute and Department of Critical Care Medicine, Maine Medical Center, Portland (P.T.M.); the Department of Pharmacy and Health Systems Sciences, Bouve College of Health Sciences, Northeastern University, Boston (J.W.D., M.S.D.); the Departments of Pharmacy (J.T.S., L.N.B.) and Surgery (J.T.S.) and the Center for Outcomes Research (J.T.S.), Houston Methodist, and the Pulmonary, Critical Care and Sleep Medicine Section, Ben Taub Hospital, Baylor College of Medicine (K.K.G.), Houston; the Division of Pulmonary/Critical Care Medicine, University of Texas Health, and the South Texas Veterans Health Care System, San Antonio (A.A.); and Texas Health Harris Methodist Hospital Fort Worth, Fort Worth (A.S.) all in Texas; the University of Sydney, and the Department of Anaesthetics, Royal Prince Alfred Hospital, Sydney, and the Department of Anesthesiology, University of Wisconsin, Madison (R.D.S.); Pulmonary and Critical Care Medicine, Baton Rouge General Medical Center and Our Lady of the Lake Regional Medical Center, Baton Rouge, LA (H.R.O.); the Department of Anesthesia and Perioperative Care, University of California, San Francisco, San Francisco (M.A.G.); Pulmonary and Critical Care, Mission Hospital, Asheville, NC (G.J.S.); the Division of Pulmonary, Critical Care and Sleep Medicine, Department of Internal Medicine, Ohio State University Wexner Medical Center, Columbus (N.E.B.); and Clinical Research, Investigation, and Systems Modeling of Acute Illness Center, Department of Critical Care Medicine, University of Pittsburgh School of Medicine, Pittsburgh (T.D.G.).

## REFERENCES

1. Fleischmann C, Scherag A, Adhikari NK, et al. Assessment of global incidence and mortality of hospital-treated sepsis: current estimates and limitations. *Am J Respir Crit Care Med* 2016;193:259-72.
2. Dhital R, Basnet S, Poudel DR. Predictors and outcome of invasive mechanical ventilation in hospitalized patients with sepsis: data from National Inpatient Sample. *J Community Hosp Intern Med Perspect* 2018;8:49-52.
3. Vincent J-L, Marshall JC, Namendys-Silva SA, et al. Assessment of the worldwide burden of critical illness: the Intensive Care Over Nations (ICON) audit. *Lancet Respir Med* 2014;2:380-6.
4. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet* 2014; 383:911-22.
5. Ely EW, Shintani A, Truman B, et al. Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA* 2004;291:1753-62.
6. Shehabi Y, Riker RR, Bokesch PM, Wisemandle W, Shintani A, Ely EW. Delirium duration and mortality in lightly sedated, mechanically ventilated intensive care patients. *Crit Care Med* 2010;38: 2311-8.
7. Pisani MA, Kong SYJ, Kasl SV, Murphy TE, Araujo KLB, Van Ness PH. Days of delirium are associated with 1-year mortality in an older intensive care unit population. *Am J Respir Crit Care Med* 2009;180:1092-7.
8. Schuler A, Wulf DA, Lu Y, et al. The impact of acute organ dysfunction on long-term survival in sepsis. *Crit Care Med* 2018; 46:843-9.
9. Pandharipande PP, Girard TD, Jackson JC, et al. Long-term cognitive impairment after critical illness. *N Engl J Med* 2013;369:1306-16.
10. Saczynski JS, Marcantonio ER, Quach L, et al. Cognitive trajectories after post-operative delirium. *N Engl J Med* 2012; 367:30-9.
11. Helmy SA, Al-Attiah RJ. The immunomodulatory effects of prolonged intravenous infusion of propofol versus midazolam in critically ill surgical patients. *Anaesthesia* 2001;56:4-8.
12. Wang X, Cheng Y, Liu X, Yang J, Munoz D, Zhang C. Unexpected pro-injury effect of propofol on vascular smooth muscle cells with increased oxidative stress. *Crit Care Med* 2011;39:738-45.
13. Kelbel I, Koch T, Weber A, Schiefer HG, van Ackern K, Neuhofer H. Alterations of bacterial clearance induced by propofol. *Acta Anaesthesiol Scand* 1999;43:71-6.
14. Memiş D, Hekimoğlu S, Vatan I, Yandim T, Yksel M, S t N. Effects of midazolam and dexmedetomidine on inflammatory responses and gastric intramucosal pH to sepsis, in critically ill patients. *Br J Anaesth* 2007;98:550-2.
15. Venn RM, Bryant A, Hall GM, Grounds RM. Effects of dexmedetomidine on adrenocortical function, and the cardiovascular, endocrine and inflammatory responses in post-operative patients needing sedation in the intensive care unit. *Br J Anaesth* 2001; 86:650-6.
16. Qiao H, Sanders RD, Ma D, Wu X, Maze M. Sedation improves early outcome in severely septic Sprague Dawley rats. *Crit Care* 2009;13:R136.
17. Nishina K, Akamatsu H, Mikawa K, et al. The effects of clonidine and dexmedetomidine on human neutrophil functions. *Anesth Analg* 1999;88:452-8.
18. Pandharipande PP, Pun BT, Herr DL, et al. Effect of sedation with dexmedetomidine vs lorazepam on acute brain dysfunction in mechanically ventilated patients: the MENDS randomized controlled trial. *JAMA* 2007;298:2644-53.
19. Riker RR, Shehabi Y, Bokesch PM, et al. Dexmedetomidine vs midazolam for sedation of critically ill patients: a randomized trial. *JAMA* 2009;301:489-99.
20. Pandharipande PP, Sanders RD, Girard TD, et al. Effect of dexmedetomidine versus lorazepam on outcome in patients with sepsis: an a priori-designed analysis of the MENDS randomized controlled trial. *Crit Care* 2010;14:R38.
21. Jakob SM, Ruokonen E, Grounds RM, et al. Dexmedetomidine vs midazolam or propofol for sedation during prolonged mechanical ventilation: two randomized controlled trials. *JAMA* 2012;307:1151-60.
22. Kawazoe Y, Miyamoto K, Morimoto T, et al. Effect of dexmedetomidine on mortality and ventilator-free days in patients requiring mechanical ventilation with sepsis: a randomized clinical trial. *JAMA* 2017;317:1321-8.
23. Shehabi Y, Howe BD, Bellomo R, et al. Early sedation with dexmedetomidine in critically ill patients. *N Engl J Med* 2019; 380:2506-17.
24. Devlin JW, Skrobik Y, Glinas C, et al. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med* 2018;46(9):e825-e873.
25. Chandrasekhar R, Hughes CG, Pun BT, Orun OM, Ely EW, Pandharipande PP. Statistical analysis plan for the Maximizing the Efficacy of Sedation and Reducing Neurological Dysfunction and Mortality in Septic Patients with Acute Respiratory Failure trial. *Crit Care Resusc* 2020;22:63-71.
26. Sessler CN, Gosnell MS, Grap MJ, et al. The Richmond Agitation-Sedation Scale: validity and reliability in adult intensive care unit patients. *Am J Respir Crit Care Med* 2002;166:1338-44.
27. Rhodes A, Evans LE, Alhazzani W, et al. Surviving Sepsis Campaign: international guidelines for management of sepsis and septic shock: 2016. *Crit Care Med* 2017;45: 486-552.
28. Barnes-Daly MA, Phillips G, Ely EW. Improving hospital survival and reducing brain dysfunction at seven California community hospitals: implementing PAD

- guidelines via the ABCDEF bundle in 6,064 patients. *Crit Care Med* 2017;45:171-8.
29. Pun BT, Balas MC, Barnes-Daly MA, et al. Caring for critically ill patients with the ABCDEF bundle: results of the ICU Liberation Collaborative in over 15,000 adults. *Crit Care Med* 2019;47:3-14.
30. Ely EW, Inouye SK, Bernard GR, et al. Delirium in mechanically ventilated patients: validity and reliability of the confusion assessment method for the intensive care unit (CAM-ICU). *JAMA* 2001;286:2703-10.
31. Góvilas C, Fillion L, Puntillo KA, Viens C, Fortier M. Validation of the Critical-Care Pain Observation Tool in adult patients. *Am J Crit Care* 2006;15:420-7.
32. Brandt J, Spencer M, Folstein MF. The telephone interview for cognitive status. *Neuropsychiatry Neuropsychol Behav Neurol* 1988;1:111-7.
33. Christie JD, Plotkin Biester RC, Taichman DB, et al. Formation and validation of a telephone battery to assess cognitive function in acute respiratory distress syndrome survivors. *J Crit Care* 2006;21:125-32.
34. Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of illness in the aged—the Index of Adl: a standardized measure of biological and psychosocial function. *JAMA* 1963;185:914-9.
35. Pfeffer RJ, Kurosaki TT, Harrah CH Jr, Chance JM, Filos S. Measurement of functional activities in older adults in the community. *J Gerontol* 1982;37:323-9.
36. Rabin R, de Charro F. EQ-5D: a measure of health status from the EuroQol Group. *Ann Med* 2001;33:337-43.
37. Jorm AF. A short form of the Informant Questionnaire on Cognitive Decline in the Elderly (IQCODE): development and cross-validation. *Psychol Med* 1994;24:145-53.
38. Strøm T, Martinussen T, Toft P. A protocol of no sedation for critically ill patients receiving mechanical ventilation: a randomised trial. *Lancet* 2010;375:475-80.
39. Olsen HT, Nedergaard HK, Strøm T, et al. Nonsedation or light sedation in critically ill, mechanically ventilated patients. *N Engl J Med* 2020;382:1103-11.
40. Djaiani G, Silverton N, Fedorko L, et al. Dexmedetomidine versus propofol sedation reduces delirium after cardiac surgery: a randomized controlled trial. *Anesthesiology* 2016;124:362-8.

Copyright © 2021 Massachusetts Medical Society.

RECEIVE IMMEDIATE NOTIFICATION WHEN AN ARTICLE  
IS PUBLISHED ONLINE FIRST

To be notified by email when *Journal* articles  
are published online first, sign up at NEJM.org.